



MiCorps Volunteer Stream Monitoring Program:
Monitoring Procedures

Updated December 2020



Prepared by:

Paul Steen, MiCorps Staff, Huron River Watershed Council
Jo Latimore, MiCorps Staff, Michigan State University

Adapted from:
“Stream Crossing Watershed Survey Procedure, April 27, 2000”

Prepared by:
Charlie Bauer, Saginaw Bay District
Greg Goudy, Cadillac District
Scott Hanshue, Great Lakes and Environmental Assessment Section Gary
Kohlhepp, Great Lakes and Environmental Assessment Section Megan McMahon,
Shiawassee District
Ralph Reznick, Nonpoint Source Unit

Surface Water Quality Division
Michigan Department of Environmental Quality

I. Overview

A. OBJECTIVES

This set of stream monitoring forms is intended to be used as a quick screening tool to increase the amount of information available on the ecological quality of Michigan's streams and rivers, and the sources of degradation to the rivers. This document is designed to provide standardized assessment and data recording procedures that can be used by trained volunteers participating in the Michigan Clean Water Corps (MiCorps) Volunteer Stream Monitoring Program.

This stream monitoring procedure is designed to address several general objectives:

- Increase the information available on the ecological quality of Michigan rivers and the sources of pollutants, for use by state biologists, local communities, and monitoring groups.
- Provide consistent data collection and management statewide.
- Serve as a screening tool to identify issues and the need for more thorough investigations.

B. TRAINING

All MiCorps Volunteer Stream Monitoring Program leaders must have received basic training in the stream assessment methods described below from MiCorps staff. Trained program leaders are then qualified to train their own volunteers in these procedures.

C. GENERAL CONCEPTS

The procedures and data forms provided below include two types of assessment: Stream Habitat Assessment and Macroinvertebrate Sampling.

The Stream Habitat Assessment is a visual assessment of stream conditions and watershed characteristics. The assessment should include approximately 300 feet of stream length. Only observations that are actually seen are to be recorded. No "educated guesses" are to be made about what should be there or is probably there. If something cannot be seen, it should not be recorded. The one exception is if a significant pollutant source or stream impact is known to be upstream of a particular site, a comment about its presence can be made in the comment section of the form.

The Macroinvertebrate Sampling procedure should be used in conjunction with the Stream Habitat Assessment because each approach provides a different piece of the stream condition puzzle. Because of their varying tolerances to physical and chemical conditions, macroinvertebrates indicate the ecological condition of the stream, while the

habitat assessment provides clues to the causes of stream degradation

Macroinvertebrate data used to calculate the Water Quality Rating (WQR), which provides a straightforward summary of stream conditions and can be used to compare conditions between study sites.

D. SURVEY DESIGN

1. Selecting Monitoring Sites

One of the basic questions in planning stream monitoring is the location of study sites: how many stream sites should be surveyed within a watershed to adequately characterize it, and where should they be located? That depends on a variety of factors including the heterogeneity of land use, soils, topography, hydrology, and other characteristics within the watershed. Consequently, this question can only be answered on a watershed-by-watershed basis.

A general EGLE guideline is to try to survey a 30% of the stream road-crossing sites within a watershed, with the sites distributed such that each subwatershed (and in turn their subwatersheds) are assessed to provide a representative depiction of conditions found throughout the watershed. At least one site should be surveyed in each tributary, with the location of this site being near the mouth of the tributary. The distribution of sampling stations within the watershed should also achieve adequate geographic coverage. Consider establishing stations upstream and downstream of suspected pollutant source areas, or major changes in land use, topography, soil types, water quality, and stream hydrology (flow volume, velocity or sinuosity). If the intent of monitoring is to meet additional, watershed-specific objectives, then additional data may be needed.

When beginning a MiCorps monitoring program, it is likely not possible to get to 30% coverage of stream road-crossing sites due to volunteer numbers and budget constraints. MiCorps will require at least 6 sites to qualify for receiving a grant. Place these as close to the mouth of different tributaries as you can, with at least two on the main branch of your system, if you have one, on public land or land you have permission to access. As your program grows, you can growth your monitoring reach to new locations.

In all cases, the site should be representative of the area of stream surveyed, it should contain a diverse range of the available in-stream cover, and it should contain some gravel/cobble bottom substrates if possible. Remember that each study site should allow for the assessment of 300 feet of stream length.

2. Time of Year and Monitoring Frequency

The time of year in which monitoring is conducted is important. For comparisons of monitoring data from year to year, data should be collected during the same season(s) each year. Ideally, macroinvertebrate sampling should take place in spring and again in early fall. Different macroinvertebrate communities are likely to be encountered during these different seasons, and sampling twice a year will provide a more complete picture of the total stream

community. All sampling must be conducted within a two-week window, and preferably, all on the same day. To provide comparable results from year to year, sampling should be conducted at approximately the same time each year.

Habitat Assessment should be done in early spring before leaf-out, or in the fall after streamside vegetation dies back, allowing visual assessments of stream characteristics. Stream habitat assessments should not be conducted when there is snow on the ground or ice on the water because important features may be hidden from view. Surveys conducted during or shortly after storm runoff events may help to identify sources of pollutants, but high-water obscures bank conditions and increased stream turbidity may make assessment of instream conditions difficult. Furthermore, all sites within a single watershed should be surveyed as closely together in time as possible to facilitate relative data comparisons among stations surveyed under similar stream flow and seasonal conditions.

MiCorps recommends repeating habitat assessment every 1 to 5 years, depending on the level of your concern for changes or impacts.

II. Stream Habitat Assessments

A. GENERAL INSTRUCTIONS

With your team (3-5 members preferably, though it can be done with 2 people), slowly walk the length of the 300 foot station length, taking in the stream's features as you go. It will be helpful to have each member be familiar with the datasheet ahead of time, so that the team knows what to look for. After observing the creek, start answering the questions together, with one member reading the questions and the other team members giving their opinions. The datasheet is filled out in a democratic method, attempting to come to agreement on the answer. If a majority agreement can't be reached, record both answers on your datasheet or where appropriate, take an average result.

Always take photos while conducting the Stream Habitat Assessment. Photographs are useful for interpretation of Stream Habitat Assessment data and for later comparisons among different sites. Site photos should show the bank conditions and some of the riparian corridor. Additional photos may be taken to highlight a particular item of concern in the stream or upland landscape. Be sure to document photos as they are taken, to simplify identification later.

As the team walks and afterwards fills out the assessment, one team member is in charge of drawing a site sketch (there is no MiCorps template for this; you can choose your methodology). The goal of a site sketch is to make the location understandable for anyone who has never been there, to make it easier to plan future outings, and to track long term changes. Draw a bird's eye view of the study site. It is important to include a north arrow, the direction of water flow, both sides of the stream channel, upland areas, parking location, and roads in the sketch, if applicable.

B. DATA SHEET

1. Stream, Team, Location Information

MiCorps Site ID#: You should create a unique numbering system for your sites. A suggested approach would be to use your organizations abbreviations and combine it with a

number. For example, HRWC-1. You want to pick a numbering system that won't accidentally copy another organization's numbering system. MiCorps staff will contact you if your numbering system is not unique.

Date: Record the month, day and year.

Time: Record the time when the monitoring activity began.

Site Name: Use a combination of the stream name and location from which you access the study site. For example, Arms Creek at Walsh Road.

Stream name: Use the stream or river name found on the U.S. Geological Survey (USGS) topographic map for the area and note also the local name if it is different. For tributary streams to major rivers, record the tributary stream name here, not the major river name. If the tributary is an unnamed tributary, record as "Unnamed Tributary to" followed by the name of the next named stream downstream. For example, a station on an unnamed tributary of Hogg Creek would be recorded as "Unnamed Tributary to Hogg Creek".

Location: This is often the name of the road from which you access the study site, or name of the public park. It is very important to indicate whether the site is upstream or downstream of the road. If the same road crosses a single stream two or more times, it is sometimes desirable to record the road name relative to the nearest crossroads (e.g. "Green Road between Brown Road and Hill Road").

Location Information: Record the latitude and longitude coordinates of the study site. Ideally, these coordinates will correspond to the midpoint of the stream study reach. Google Maps now allows for very easy latitude/longitude identification. Just right click on the map and these coordinates will be given.

Names of Team members: Record the name of all the team members participating in the assessment, and circle the one recording the data, in case questions come up later.

2. Stream and Riparian Habitat

A. General Information

1. Avg. Stream Width (ft): Circle the range that represents the average stream width in feet.

This can be a best guess, or you can choose to take width measurements of the stream at several points along the 300-foot assessment area, and indicate the average width here. These measurements are also useful in creating the Stream Site Sketch.

2. Avg. Stream Depth (ft): Circle the appropriate depth range in feet. Take depth measurements at several points within the 300-foot assessment area and take the average depth. This observation is for the average depth of the stream that is consistently observed. For example, if the stream is generally shallow (<1ft), but has a pool that is 3ft deep, circle the <1ft category since a pool is not representative of the average depth of <1ft observed over most of the stream.

3. Has this stream been channelized? Stream shape constrained through human activity- look for signs of dredging, armored banks, straightened channels.

Yes, currently: You see active construction, or vegetation removal, or scraping of banks, and the river lacks turns and meanders.

Yes, sometimes in the past: The river lacks turns and meanders, but there are signs of water flow induced erosion, and vegetation has recovered from any construction at the site.

No: The stream has bends and meanders and you do not see the signs noted above. (note that you might only notice bends and meanders in small creeks; rivers bend and meander at a much higher geographic scale)

4. Estimate of current stream flow: All of these pieces of information can help you make this determination. 1) The volunteers knowledge of recent weather conditions (e.g. how much it has rained recently). 2) Visual stream observations (look for event related conditions water running off the land into the stream, fast stream water velocity, increased water turbidity, an increase in the amount of debris being carried by the stream), 3) The teams knowledge (or best guess) of what is typical flow for that (or a similar) stream, in that geographic area, for that season of the year.

Dry = No standing or flowing water, sediments may be wet.

Stagnant = Water present but not flowing, can be shallow or deep.

Low = Flowing water present, but flow volume would be considered to be below average for the stream.

Medium = Water flow is in average range for the stream.

High = Water flow is above average for the stream.

5. Highest water mark (in feet above the current level): Look for signs that the water was once higher: debris trapped against bridges, or trees, and erosion along banks above the water level.

6. Which of these habitat types are present?

Good quality streams have a wide variety of habitat available to fish and macroinvertebrates to: (1) protect them from predators, (2) avoid certain stream conditions such as fast flow velocities or direct sunlight, and 3) provide surfaces and structure on which food grows, collects, or tries to hide. Circle all the habitat types on the data form that are present in the stream reach for your 300 foot station. Types of habitat include the following:

Riffles: Riffles are areas of naturally occurring, short, relatively shallow, zones of fast moving water, typically followed by a pool. The water surface is visibly broken (often by small standing waves) and the river bottom is normally made up of gravel, rubble and/or boulders. Riffles are not normally visible at high water and may be difficult to identify in large rivers. The size of, and distance between, riffles is related to stream size. In large mainstream reaches, such as the Manistee or Muskegon rivers, riffles may be present. in the form of rapids.

Pool: Pools are areas of relatively deep, slow moving water. The key word here is "relatively". Water depth sufficient to classify an area as a pool can vary from around 8 inches in small streams, to several feet in wadable streams, to tens of feet in large rivers. Pools are often located on the outside bend of a river channel and downstream of a riffle zone or obstruction. The water surface of a pool is relatively flat and unbroken. The presence of pools in large rivers may be difficult to identify because of an increase in relative scale, and an often-limited ability to see to the bottom of deep or turbid stream reaches.

Large woody debris: Logs, branches, and roots both above and below the water surface.

Large rocks: rocks that are 10 inches in diameter or larger.

Undercut Banks: Stream banks that overhang the stream because water has eroded some of the material beneath them.

Overhanging Vegetation: Terrestrial vegetation that extends out from shore over the surface of the stream within a foot or two of the water surface (includes trees, shrubs, grasses, etc.). This category also includes sweeping vegetation, which is terrestrial shoreline vegetation that extends into the water itself (such as low hanging branches on shrubs) and is therefore often “swept” in a downstream direction by the current.

Rooted Aquatic Plants: Aquatic macrophytes provide breaks in water flow, cover, and a food source, becoming good habitat for both fish and macroinvertebrates.

7. Estimate of turbidity: Water appears cloudy—it is rarely transparent, and the level of the cloudiness is called turbidity. Turbidity is caused by suspended particulates such as silt, sand, algae, or fine organic matter. Highly turbid water is opaque to varying degrees, preventing the observer from seeing very far into it. Note that water can have a color to it that is not turbidity, such as the brown transparent water often associated with swampy areas.

8. Is there a sheen or oil slick visible on the surface of the water?

9. If yes to #8, does the sheen break up when poked with a stick?

An oily appearing sheen on the water surface caused by petroleum products. A thin sheen will often have a rainbow of hues visible. The sheen can be distinguished from bacterial sheens by remaining viscous when poked with a stick or otherwise physically disturbed, whereas bacterial sheens break into distinct platelets.

10. Is there foam present on the surface of the water?

11. If yes to #10, does the foam smell soapy and look white and pillow like or look gritty with dirt mixed in?

Naturally occurring foam often looks like soap suds on the water surface and can be white, grayish or brownish. Foam is produced when water with dissolved organic material is aerated and can range in extent from individual bubbles to mats several feet high. Foam is typically produced in streams when water flows through rapids or past surface obstructions such as logs, sticks and rocks. Simple wave action can produce foam in lakes. This naturally occurring foam is quite common. If the suds are a bright white color, billowy and pillow-like, soapy, or smell perfumed, it is not natural foam. Volunteers used to touch the foam to feel for grittiness, but MiCorps does not advise that anymore as the foam could be PFAS, which you should not handle.

The following are optional measurements not currently funded by MiCorps (water temperature, dissolved oxygen, pH, water velocity)

B. Streambed Substrate

Substrate is the material that makes up the bottom of the stream. In general, good quality substrates (from an aquatic habitat perspective) contain a large amount of coarse aggregate material—such as gravels and cobbles—with a minimal amount of fine particles surrounding or covering the interstitial pore spaces. These stable materials provide the solid surfaces necessary for the colonization of attached algae and the development of diverse macroinvertebrate communities.

Using the particle size and composition guidance provided below, identify the percent areal extent of each substrate type present. The composition estimate should include the entire area of the stream bottom in the study site (typically, 300 feet of stream). Sometimes it is not possible to determine the substrate type all the way across a river because it is too deep or the water is turbid. In these cases, assign the appropriate percentage amount to the “unknown” category.

Substrate Type and Sizes

Boulder: Rocks 10 inches diameter or larger.

Cobble: Rocks 2.5 inch to 10 inches in diameter.

Gravel: 0.1 -2.5 inch diameter

Sand: Coarse grained, <.1 inch diameter particles

Silt-Muck-Detritus: Silt is usually clay, very fine sands, or organic soils, 0.004 to 0.06 millimeters in diameter. Muck is decomposing organic material of very fine diameter. Detritus is small particles of organic material such as pieces of leaves, sticks, and plants.

Hardpan-Bedrock: Solid surface. Hardpan is usually packed clay. Bedrock is a solid rock surface (the tops of buried boulders are not bedrock).

Artificial: Human made, such as concrete piers, sheet piling or rock riprap (that portion of shoreline erosion protection structures that extends below the water surface is considered substrate).

Other (specify): If something doesn't fit into the above categorizes, it goes here.

Can't see: The portion of the stream bottom for which a substrate type determination cannot be made because the bottom cannot be seen due to water depth or turbidity.

C. Bank stability and erosion

Bank erosion may occur as a result of natural flow conditions, or may be caused by human activities. Determine the severity of erosion that has taken place through the

explanations given for the categories excellent, good, marginal, and poor, and then circle one of the numbers in that category to give a more specific rating.

Excellent: Banks Stable. No evidence of erosion or bank failure. Little potential for problems during floods. < 5% of bank affected.

Good: Moderately stable. Small areas of erosion. Slight potential for problems in extreme floods. 5-30% of bank in reach has areas of erosion.

Marginal: Moderately unstable. Erosional areas occur frequently and are somewhat large. High erosion potential during floods. 30-60% of banks in reach are eroded

Poor: Unstable. Many eroded areas. > 60% banks eroded. Raw areas frequent along straight sections and bends. Bank sloughing obvious.

D. Plant Community

Estimate the percentage of the stream covered overhanging vegetation/tree canopy? Circle one: <10%, 10-50%, 50-90%, >90%. These are very wide windows because a general sense of the situation is all that is needed. Is the stream fully exposed to the sun, fully shaded, or somewhere in between? The level of sun exposure will affect how biota hides and water temperature fluctuations.

For the various type of plants listed, rate each group as absent, rare, common, or abundant. The groups are:

Plants in the Stream:

Floating Algae: The abundance of suspended algae (single celled organisms that may or may not form colonies) or algae on the surface or rocks or plants should be recorded here.

Filamentous Algae: Algae that appear in stringy or ropy strands, such as Cladophora. The strands may or may not be attached to other objects in the waterbody.

Macrophytes: This category refers to aquatic plants. By definition, macrophytes are any plant species that can be readily seen without the use of optical magnification. However, the usage here is directed primarily toward aquatic vascular plants—plants with a vascular system that typically includes roots, stems and/or leaves. This includes duckweed, as it is a floating vascular plant. Certain large algae species that superficially look like vascular plants, such as Chara, can be recorded here as well. If the person conducting the survey is knowledgeable about aquatic plants, the particular type or species of plant(s) can be noted in the comment section at the end of the form. Floating, suspended, or filamentous algae species should be recorded in one of the algae categories and not here.

Plants on the bank/riparian zone

Shrubs: Woody, low lying plants.

Trees: Woody, tall plants.

Herbaceous: Non-woody plants including grasses, forbs, and so on.

E. Riparian Zone

The riparian vegetative width is the width of the streamside natural vegetation zone along the stream banks. The width is measured from the edge of the stream to the end of the contiguous block of natural vegetation. Natural vegetation is defined as including trees, shrubs, old fields, wetlands, or planted vegetative buffer strips (often used in agricultural areas and stormwater runoff control). Agricultural crop land and lawns are not considered natural vegetation for the purposes of this question. A stream with grass mowed to the very edge is said to have no riparian zones. A stream set in a deep forest will have a riparian zone that spreads further than you can even see.

For both the left and right bank (which is determined by looking downstream), circle the landuse types that you can see along your 300 foot stretch.

Then, rate the riparian zone from excellent to poor, and then circle one of the numbers in that category to give a more specific rating, similar to how you rated bank erosion in C.

Excellent: Width of riparian zone >150 feet, dominated by vegetation, including trees, understory shrubs, or non-woody macrophytes or wetlands; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.

Good: Width of riparian zone 75-150 feet; human activities have impacted zone only minimally.

Marginal: Width of riparian zone 10-75 feet; human activities have impacted zone a great deal.

Poor Width of riparian zone ,10 feet; little or no riparian vegetation due to human activities.

III. Sources of Degradation

The intent of this section is to evaluate the relative importance of potential sources in terms of pollutant contribution to the waterbody at a given site in the watershed. The evaluation assesses the potential for pollutant inputs at the site, NOT pollutant impacts, or the potential for pollutant impacts. Pollutant impacts, as indicated by visual manifestations (like erosion, changes to substrate, oil, foam, etc) were evaluated previously.

Evaluating potential sources of pollutants to a waterbody is a three step process: identification of potential sources, evaluation of pathways for pollutants to get to the waterbody, and finally evaluation of the severity (magnitude) of this pollutant input or loading. The three steps of this process will result in scoring identified sources on the survey sheet as Slight, Moderate, or High Priority in terms of the severity or amount of their pollutant contribution to the waterbody at the site being surveyed.

(1) Source Identification

Visually evaluate the various land use/land change activities at the site for potential sources of pollution. Note all potential sources for the area that can be seen (choosing from among the list of sources on the data sheet). For example, is there evidence of soil disturbance at the site, or land uses such as residential lawns, agricultural fields, parking lots, urban areas, etc., near the waterbody? Use the source definitions provided to help identify what potential sources may exist. If it is known that a significant source exists upstream of the study site, such as a wastewater treatment plant, it may be important to note the presence of that source, but it should be recorded in the comments section since it was not visible at the site.

(2) Pollutant Pathway

Next, for each potential source that has been identified, evaluate how pollutants could get from the source to the water. An evaluation of likely pathways for pollutants to enter the waterbody provides information regarding the potential for the identified sources to contribute pollutants. The following provides a quick outline of some visual observations to consider in evaluating pollutant pathways. Pay particular attention to likely water runoff patterns at the site that may occur during rainfall or snowmelt events.

- Gully/rill erosion provides a direct pathway for pollutants to enter the stream in a concentrated flow when the land slopes toward the stream. Pollutants associated with eroding soils will vary depending on the type of land use activity.
- Tile/pipe discharges are potential direct pathways for pollutants.
- Bare soils near the edge of a waterbody provide a likely pathway for sediment to get to the waterbody.
- Maintained lawns to the edge of a waterbody provide a likely pathway for nutrients and pesticides to the waterbody.
- Land disturbance/use activities to the edge of a waterbody provide a likely pathway for various pollutants to the waterbody.
- Open areas of disturbed soils and/or bare soils devoid of vegetation provide a potential pathway for pollutants via wind erosion.
- Steep streambanks (steeper than a 2:1 slope) devoid of vegetation are likely pathways for sediment.
- No canopy over the waterbody is a pathway for dramatic thermal increase in water temperature during the day.
- Impervious surfaces (parking lots, roads, roof tops, etc.) provide a likely pathway for various pollutants, and may increase flows in the watershed causing flashiness.

- Culverts/bridges may not be aligned with the stream, or may be undersized, and could provide a likely pathway for flow to create streambank erosion both upstream and downstream of the culvert or bridge.

(3) Severity Ranking

Finally, for each source for which a pathway has been identified, evaluate how severe the pollutant loading is. Rank each source identified as Slight, Moderate or High severity for the contribution of pollutants, based on the magnitude or quantity of pollutants likely to be delivered to the stream. The surveyor must use their judgement on assigning a slight, moderate, or high rating.

The severity ranking is based only on pollutant inputs from the specific source at the site, not on visible stream impacts or impacts the pollutant may cause downstream. The pollutant loads from the identified source(s) may or may not have an impact at the site.

Evaluation of the source, location and pathways can provide a reasonable assessment of the severity of the pollutant loading. The following provides a quick outline of some visual observations to consider in evaluating the severity of pollutant loading.

- Proximity to waterbody – generally the closer the use, or land disturbance activity, is to the waterbody, the greater the likelihood for pollutant delivery.
- Slope to waterbody – generally the steeper the slope/topography to the waterbody, the greater the likelihood of overland pollutant delivery.
- Conveyance to waterbody (ditch, pipe, etc.) – generally a conveyance from the use, or land disturbance activity, increases the likelihood of pollutant delivery.
- Imperviousness – impermeable surfaces reduce the amount of land area available for water infiltration and increase the potential for overland runoff. Additionally, if a watershed is greater than 10% impervious, it will start to show some systemic problems due to impacts from flow. If a watershed is greater than 25% impervious, the natural hydrology is generally heavily impaired.
- Intensity and type of use, or land disturbance activity – generally the more intensive the activity the greater the likelihood for the generation of pollutants. Certain activities may have specific types of pollutants associated with them.
- Size of erosion area – generally the larger the erosion area the greater the likelihood for sediment delivery.
- Soil type – clay is less permeable than sand, and therefore would create a greater potential for overland runoff of pollutants.
- Presence and type of vegetation – the greater the vegetative buffer around a waterbody, the better the filtration of pollutants from nearby land disturbance and use activities. Certain types of vegetative buffers work better than others and should be evaluated on a case-by-case basis.

Potential Source Category Definitions:

Source Category	Use this Source Category if ...
Crop Related Sources	... there is a reasonably clear pathway for pollutants to enter the waterbody from the farmed area. Possible pathways: farming to the edge of the drain, gully/rill erosion off field, tile discharge, wind erosion off field.
Grazing Related Sources	... there is clear evidence that grazing of animals near or in the waterbody has resulted in the degradation of streambanks or stream beds, sedimentation, nutrient enrichment, and/or potential bacterial contamination.
Intensive Animal Feeding Operations	... there is a reasonably clear pathway for pollutants to enter the waterbody from either runoff from the operation or land application of animal manure. Possible pathways: overland flow, tile discharge.
Highway/Road/Bridge Maintenance and Runoff (Transportation NPS)	... there is clear evidence that transportation infrastructure is creating increased flow, runoff of pollutants, or erosion areas in or adjacent to the waterbody.
Channelization	... there is clear evidence that the natural river channel has been straightened to facilitate drainage.
Dredging	... there is clear evidence that a waterbody has been recently dredged. Evidence might include: spoil piles on side of waterbody, disturbed bottom, disturbed banks.
Removal of Riparian Vegetation	... there is clear evidence that vegetation along the waterbody has been recently removed (within the last few years).
Bank and Shoreline Erosion/Modification/Destruction	... there is clear evidence that the banks or shoreline of a waterbody have been modified through either through human activities or natural erosion processes.
Flow Regulation/Modification (Hydrology)	... there is reasonably clear evidence that flow modifications in the watershed have created unstable flows resulting in streambank erosion.
Upstream Impoundment	... there is reasonably clear evidence that an upstream impoundment has contributed to impacts on downstream sites. Impacts may be: nuisance algae, increased temperatures, streambank erosion from unstable flows.
<u>Construction:</u> Highway/Road /Bridge/Culvert	... there is clear evidence that on-going or recent construction of transportation infrastructure is contributing pollutants to the waterbody.
<u>Construction:</u> Land Development	... there is clear evidence that on-going or recent land development is contributing pollutants to the waterbody.
Urban Runoff (Residential/ Urban NPS)	... there is a reasonably clear pathway for pollutants to enter the waterbody from an urban/residential area. Possible pathways: gully/rill erosion, pipe/storm sewer discharge, wind erosion, runoff from lawns or impervious surfaces.
Land Disposal	... there is a reasonably clear pathway for pollutants to enter the waterbody from an area where waste materials (trash, septage, hazardous waste, etc.) have been either land applied or dumped. Possible pathways: gully/rill erosion, pipe discharge, wind erosion, or direct runoff.

On-site Wastewater Systems (e.g. septic systems)	... there is reasonably clear evidence of nutrient enrichment and/or sewage odor is present, and there is reason to believe the area is unsewered.
Silviculture (Forestry NPS)	... there is a reasonably clear pathway for pollutants to enter the waterbody from the forest management area. Possible pathways: logging to the edge of the waterbody, gully/rill erosion off site, pumped drainage, erosion from logging roads, wind erosion off site.
Resource Extraction (Mining NPS)	... there is a reasonably clear pathway for pollutants to enter the waterbody from the mined area. Possible pathways: gully/rill erosion off site, pumped drainage, runoff from mine tailings, wind erosion off site.
Recreational/Tourism Activities (general)	... you are unable to clearly identify the recreational source as related to a golf course, or recreational boating activity. Foot traffic causing erosion would fall into this category.
Golf Courses	... there is a reasonably clear pathway for pollutants to enter the waterbody from the golf course area. Possible pathways: overland runoff, gully/rill erosion off course, tile discharge, wind erosion off course.
Marinas/Recr. Boating (water releases)	... if you can reasonably determine that releases of pollutants to a waterbody such as septage or oil/gasoline are due to recreational boating activities.
Marinas/Recr. Boating (streambank erosion)	... you can reasonably determine that streambank erosion is due to wake from recreational boating activities.
Debris in Water	... debris in the water either is discharging a potential pollutant, or is causing in stream impacts due to modifications of flow. Possible examples: Leaking barrel, Refrigerator, Tires, etc. This does not include general litter (e.g. paper products).
Industrial Point Source	... there is reasonably clear evidence that an upstream industrial point source has contributed pollutants.
Municipal Point Source	... there is reasonably clear evidence that an upstream municipal point source has contributed pollutants.
Natural Sources	... there is reasonably clear evidence that natural sources are contributing pollutants. Possible examples: streambank erosion, pollen, foam, etc.
Source(s) Unknown	... if you see an impact but are unable to clearly identify any likely sources.

Additional Comments:

Any observations about the site that were not covered elsewhere on the survey form should be recorded in this section. If certain survey responses require clarification or elaboration, those should be described here as well. The comment section can also be used to add detail to the site characterization, such as listing the types of aquatic plants or algae present, if known.

In addition, any unique conditions or issues that arose or were observed during the assessment process should be noted here.

IV. *Optional Quantitative Measurements*

A. Transects and Pebble Counts

To take quantitative stream habitat measurements, conduct 10 transects of your stream reach. A transect is a measuring tape line stretched out perpendicularly across the stream, going from bank to bank. At 10-20 locations along this line, you will take depth measurements and record the substrate type.

Required equipment: tape measure long enough to stretch across the stream, and graduated rod or stick to measure water depth. Data sheet is on the next page.

Directions:

- 1) Determine stream width.
- 2) Use the rod to measure depth (D) and substrate (S) at more than 10 but less than 20 regular intervals along the entire transect. (For streams less than 10 feet wide, measure every ½ foot, for streams about 10 feet wide, measure every foot, etc.)
- 3) At every depth measurement, identify the single piece of substrate that the rod lands on. If it is a mix of substrates, randomly pick one of them, and the next time you find a similar grouping, pick the other(s).
- 4). For every measurement, enter the reading on the tape measure, the depth, and the substrate on the data sheet on the next page.

Data use: The depth and tape measure reading can be used to produce stream cross-section profiles. The pebble count can be used to give a more accurate percentage breakdown of the stream substrate than simply making an eyeball estimate (see Section II-B).

B. Bank Height

Vertical banks higher than 3 feet are usually unstable, while banks less than 1 foot, especially with overhang, provide good habitat for fish. While doing the transects, measure bank heights and record the angle of the bank (right, acute, or obtuse) as indicated on the data sheet. Left/right banks are identified by looking downstream.

Data use: Calculate the percentage of banks with right, obtuse, and acute angles. Right angles indicate higher erosive potential, while acute angles improve the habitat structure of a stream.

V. *Final Check*

Completeness: A volunteer team member other than the person who filled out the data sheets must check the data sheet for completeness before the team leaves the site. This verification of completeness should be noted at the bottom of each page.

Name of person who entered data into data exchange: This field is for use in case problems come up with the data entry.

Date of data entry: This field is for use in case problems come up with the data entry.

STREAM TRANSECT DATASHEET


B: Boulder -- more than 10"
 C: Cobble -- 2.5 - 10"
 G: Gravel -- 0.1 - 2.5"
 S: Sand -- fine particles, gritty

F: Fines: Silt/Detritus/Muck
 H: Hardpan/Bedrock
 A: Artificial
 O: Other (specify)

T= Reading on tape
 D = Depth
 S = Substrate

Stream Width	EXAMPLE			Transect #			Transect #			Transect#		
	T	D	S	T	D	S	T	D	S	T	D	S
Beginning Water's Edge:	1.5											
1	2.5	0.4	G									
2	3.5	0.4	G									
3	4.5	0.4	G									
4	5.5	0.2	C									
5	6.5	0	S									
6	7.5	0.6	S									
7	8.5	0.7	G									
8	9.5	0.7	G									
9	10.5	0.6	C									
10	11.5	0.7	B									
11	12.5	0.4	G									
12	13.5	0.3	F									
13	14.5	0.2	F									
14												
15												
16												
17												
18												
19												
Ending Water's Edge:	14.8											
Bank Side	L	R		L	R		L	R		L	R	
Bank Height	1.7 feet	0.5 feet										
Does the bank have an undercut?	N	Y										
If so, how wide is it?		1 ft										
Bank Angles: Sketch												

Sketch examples:



Undercut
(Acute)

Obtuse

Right

III. Stream Macroinvertebrate Monitoring Protocols

A. TEAM COMPOSITION

MiCorps macroinvertebrate collection is carried out by teams of staff and/or volunteers consisting of no fewer than 3 people and up to 6 or 7. More people than that is acceptable but as more join a team, crowding and equipment issues can hamper team effectiveness.

One team member is the Collector, who must be trained in collection techniques. This person is the only one to enter the water and use the net to pull out debris and macroinvertebrates. However, on larger rivers or streams with overgrown banks it is helpful to have a Collector's Assistant in waders assisting the Collector by carrying trays back and forth from the Collector to the Pickers.

There should also be a Team Leader, who has preferably been to a special training but at a minimum has participated in the monitoring previously. The Team Leader directs the rest of the team, the Pickers, who do not have to be trained ahead of time. On-site directions are sufficient as the Picker role is very easy and done under direct supervision of the Team Leader. The Pickers and Leader sit on the bank of the stream to pick insects out of the trays and put the specimens in the sample vials. The Team Leader also fills out data sheets, watches the time, and keeps the team organized.

B. SAMPLING

The sampling effort expended to collect benthic macroinvertebrates at each 300 foot site should be sufficient to ensure that all types of benthic invertebrate habitats are sampled in the stream reach. This generally will be about 35-45 minutes of total sampling time per station. You should be flexible on the timing for Collectors who move slowly in the water, because of either tricky wading and walking conditions or inexperience. If sampling goes slow, sample longer than 45 minutes at your discretion; the goal is to keep the total effort the same across all sampling outings.

Macroinvertebrate samples should be collected from all available habitats within the stream reach using a dip net with a 1-millimeter (mm) mesh, or by hand picking bigger items like logs and rocks.

Available habitat types can include but aren't limited to riffles, pools, cobbles, aquatic plants, runs, stream margins, leaf packs, undercut banks, overhanging vegetation, and submerged wood. Habitat and substrate types from which macroinvertebrates were collected (or collections were attempted) should be recorded on the form; include as many as possible. People on the bank can aid the Collector by reminding them of the different habitat types to sample.

As the Collector obtains debris in their net, the debris is dumped into white trays along the bank. The Pickers will then sort through the debris and place the macroinvertebrates into jar(s) of 70% ethanol preservative for later identification. The Team leader should show Pickers how to sort through the tray, and to inspect rocks and other debris, emphasizing hidden locations under bark and in caddisfly cases. The Team leader should stress

patience. Use some water to get things moving as a dry sample is nearly impossible to pick through.

Be sure that every jar has a laser printer label (or handwritten with pencil) to avoid the ink running. Place labels inside the jar with the alcohol and not taped to the outside.

The Pickers should work for about one hour in total or until they have gone through all the debris provided by the Collector, whichever comes first. The team should set a timer or mark the start time in order to be accurate. The teams must strive to get at least 100 specimens. They are not expected to count it, but generally they should have a good sense as they go if they are meeting that benchmark. The Water Quality Rating (WQR) is designed to be most accurate with sample sizes of at least 100 specimens.

C. COLLECTING TECHNIQUES IN DIFFERENT HABITATS

General Techniques

1. Collecting should begin at the downstream end of the stream reach and work upstream.
2. Please note that many mussels are endangered or threatened. Don't collect mussels and clams; don't even take them out of the water or dislodge them. Make a note on the datasheet if they are found.
3. While crayfish are not endangered, they are too big usually to fit in sample jars. Make note of crayfish and then release them as well.
4. Remember - BE AGGRESSIVE- the animals are holding on tight to rocks, branches, and leaves to avoid being carried downstream and you want to shake them loose!
5. Always point opening of net upstream so the current does not wash out your net.
6. Lift up carefully in sweeping motions to avoid losing organisms.

Riffles/Runs:

1. Keep in mind that flow has a big impact on the types of animals that can live there. Both riffles and runs are areas of faster moving water. A riffle (white water present, larger rocks) and a run (no white water, smaller gravel sized rocks) will likely yield different animals.

1. Put net on bottom of stream, stand upstream, hold net handle upright.
2. Use kicking/shuffling motion with feet to dislodge rocks. You are trying to shake organisms off rocks as well as kick up organisms that are hiding under the rocks. Dig down with your toes an inch or two. Some people use their hands to rub organisms off rocks, but beware of sharp objects on the stream bottom.

Quiet Place/pool:

1. Scoop some sediment up in your net. Some animals burrow into the muck.
Tip: When your net is full of muck, it is very heavy. To clean the excess muck out of your net: keep the top of the net out of the water to avoid losing animals, then sway the net back and forth, massaging the bottom of the net with your hand. When choosing a soft bottom area try to find one that contains silt since it is a far more productive habitat than just sand.
2. Don't oversample muck. Not much will live here, and it is difficult to sort through. Process one or two nets worth and then don't go back to this habitat.

Undercut Bank/Overhanging Vegetation or Roots:

1. Jab the net into the undercut bank while pulling the net up. Move in a quick bottom to surface motion to scrape the macroinvertebrates from roots. Do this several times.
2. If you notice roots or overhanging vegetation, put the net under the bank at the base of the plants. Shake the vegetation using your net, trying to shake off the animals clinging to these plants. Feel free to use your hands if you are sure the plants are not poisonous.

Submerged or emergent vegetation:

1. Keeping the net opening pointed upstream, move the net through vegetation trying to shake the vegetation and catch any animals.
2. Use your hands to agitate the vegetation and dislodge the animals into the net.

Rocks/Logs:

1. Small logs and rocks can be pulled out of the water by hand and given to the team to search for animals.

Hint for Logs: Be sure to check under bark.

Hint for Rocks: Caddisfly homes often look like small piles of sticks, clumps of small gravel, or even tiny circular pieces of algae attached to rocks.

Leaf Packs:

1. Look for a decomposing leaf pack. A “good” leaf pack has dark brown-black skeletonized leaves. Slimy leaves are an indication that they are decaying. Scoop a few into your net and let the team pull them apart and look for animals.
2. Sometimes a little water in the pan with the leaves will help dislodge the animals.

D. CLEANING YOUR GEAR

Remember to clean the net and pans before leaving the site to avoid transporting animals or plants. If you plan to use the gear again within the next month, air drying is not sufficient. In that case, you must clean out the treads of the waders, get all dirt of debris out of the equipment, and use a dilute bleach or similar disinfectant to sanitize the gear. For full instructions on decontamination processes, see <https://www.hrwc.org/volunteer/decontaminate/>.

E. IDENTIFICATION

Identification can be performed in the field or in an indoor setting (recommended), as desired by the monitoring organizations. Volunteers who lack identification experience must be overseen by an identification expert or program’s scientific advisor; in any case, the final identification must be confirmed by this person(s).

The organisms in the collection should be identified to order, sub-order, or family, as indicated on the MiCorps datasheet, using taxonomic keys. The abundance of each taxon in the stream study site should be recorded on the datasheet.

F. STREAM MACROINVERTEBRATE DATASHEET

Front page

MiCorps Site ID#: You should create a unique numbering system for your sites. A suggested approach would be to use your organizations abbreviations and combine it with a number. For example, HRWC-1. You want to pick a numbering system that won't accidentally copy another organization's numbering system. MiCorps staff will contact you if your numbering system is not unique.

Site Name: Use a combination of the stream name and location from which you access the study site. For example, Arms Creek at Walsh Road.

Stream name: Use the stream or river name found on the U.S. Geological Survey (USGS) topographic map for the area and note also the local name if it is different. For tributary streams to major rivers, record the tributary stream name here, not the major river name. If the tributary is an unnamed tributary, record as "Unnamed Tributary to" followed by the name of the next named stream downstream. For example, a station on an unnamed tributary of Hogg Creek would be recorded as "Unnamed Tributary to Hogg Creek".

Location: This is often the name of the road from which you access the study site, or name of the public park. It is very important to indicate whether the site is upstream or downstream of the road. If the same road crosses a single stream two or more times, it is sometimes desirable to record the road name relative to the nearest crossroads (e.g. "Green Road between Brown Road and Hill Road").

Date: Record the month, day and year.

Collection Start Time: Record the time when the monitoring activity began.

Major Watershed: Record the name of the major watershed where the study site is located (e.g., Grand River Watershed, St. Mary's River Watershed), and the corresponding HUC Code, if known.

Longitude and Latitude: Record the latitude and longitude coordinates of the study site. Ideally, these coordinates will correspond to the midpoint of the stream study reach.

Names of Team members: Record the name of all the team members participating in the assessment, and circle the one recording the data, in case questions come up later.

Stream Conditions: This section is important for interpreting the data after the collection and identification. If results are much worse than normal, this information will help the program manager conclude that conditions on the sample day were not representative of the stream's normal range of conditions and may flag the site for resample or strike the results from the long-term dataset.

Average Water Depth: This value can be taken from the Stream Habitat Assessment datasheet, if completed at the same time. Otherwise, to measure average water depth (ft), three measurements should be made at random points along the representative reach length being surveyed, and these values averaged for a mean depth.

Notable weather condition of the last week: Substantial rainfall or drought especially can cause fluctuations in macroinvertebrate results.

Are there are current site conditions that may impede normal macroinvertebrate sampling? This is left open for volunteers to comment on anything that would affect the study (for example, weather, flooding, poor visibility like high turbidity, difficult wading conditions, etc).

Habitat types: A list of stream microhabitat are provided so that the Streamside Leader can remind the Collector of what different places to sample. Sample as many of these as possible, checking them off as you go.

Did you see any crayfish or clams/mussels? Do not collect these, but record the number that you see so you can use them in your water quality rating.

Collection Finish Time and Picking Finish Time: Record the time the collector stops their work in the stream and the time when Pickers put the last specimen in the collection jars.

Identifications made/supervised: Record who was responsible for giving the final identification of the specimens.

Backpage:

Identification and Assessment:

MiCorps requires stream monitoring programs to identify macroinvertebrates to the Order level primarily, sometimes sub-Orders, and sometimes Family. This system was built to be a balance between scientific accuracy and ability of volunteers to learn how to identify insects with a moderate level of effort. While requiring genus-species level identification would be most scientifically accurate, it would prevent the program from being conducted as a volunteer program.

With counts and identifications complete, it is possible to produce a single score for the site. This scoring system is based on the Hilsenhoff Biotic Index, a scheme established by Dr. William Hilsenhoff, a famous (for this field) entomology professor from the University of Wisconsin Madison. Hilsenhoff and those who took up his work afterwards have assigned pollution sensitivity ratings to most macroinvertebrate species, genera, and families. Using the sensitivity ratings, a type of weighted average can be calculated to generate the pollution tolerance rating (or water quality rating) for macroinvertebrate samples on a scale of 0 (very pollution sensitive) to 10 (very pollution tolerant).

In MiCorps protocols, we are not identifying macroinvertebrates to the lower taxonomic levels, so leeway had to be taken with Hilsenhoff's sensitivity score to produce an average sensitivity rating for each of the taxonomic groups on the datasheet. This was done by averaging the sensitivity ratings of the different families and assigning the result to the larger taxonomic group. For example, the sensitivity ratings for the eight families of stoneflies found in Michigan were averaged for a result of 1.1. Thus 1.1 is the sensitivity for MiCorps Stonefly group.

In other words, the sensitivity ratings that MiCorps uses are best estimates for that taxonomic order but are not perfect. Again, this loss of accuracy is because of the balance that needs to be met between identification and volunteer/program leader ability.

The final MiCorps score given to each site is called the WQR (Water Quality Rating).

To calculate the WQR, follow these steps:

1. As you identify your macroinvertebrates, record the number you found for each type in the left column marked "Count". When you are done, add up all the "Count" column to get a total abundance.
2. Multiply the "Count" by the given Sensitivity Rating for each taxa group and record it in the column "Count x Sensitivity". For example, if you found 30 mayflies you would multiply 30 x 3.4 and record 102 in the "Count x Sensitivity" column.
3. Add up all the values in the "Count x Sensitivity" column and record this in the box "Sum of (Count x Sensitivity)".
4. Divide the "Sum of (County x Sensitivity)" by the "Total Abundance." The result is the site's Water Quality Rating (WQR). The lower the score, the more pollution sensitive insects are found, and the better the water quality.
5. Important Note about Abundance: This rating scale does not work when macroinvertebrate abundance is low, as a few sensitive taxa can pull the score down to very healthy levels, biasing the results. To correct for this, if abundance is less than 30, the site is automatically given a WQR of 10 (very poor). If the abundance is less than 60, the site is automatically given a WQR of 7 (poor rating). Teams should be striving to collect at least 100 specimens from each site. If the team collects from 60-99 specimens, then score the site as normal and input it into the MiCorps data exchange as normal but consider the rating to be somewhat tentative and strive for higher abundances in future visits.

Water Quality Rating		Degree of Organic Pollution
0.0-3.50	excellent	Pollution unlikely
3.51-4.50	very good	Slight pollution possible
4.51-5.50	good	Some pollution possible
5.51-6.50	fair	Fairly substantial pollution likely
6.51-7.50	fairly poor	Substantial pollution likely
7.51-8.50	poor	Very substantial pollution likely
8.51-10.0	very poor	Severe pollution likely