

# 2023 Data Report for

# Second Fortune Lake, Iron County

Site ID: 360177

# 46.0772°N, 88.4265°W

The CLMP is brought to you by:



#### About this report:

This report is a summary of the data that have been collected through the Cooperative Lakes Monitoring Program. The contents have been customized for your lake. The first page is a summary of the Trophic Status Indicators of your lake (Secchi Disk Transparency, Chlorophyll-a, Spring Total Phosphorus, and Summer Total Phosphorus). Where data are available, they have been summarized for the most recent field season, five years prior to the most recent field season, and since the first year your lake has been enrolled in the program.

If you did not take 8 or more Secchi disk measurements or 4 or more chlorophyll measurements, there will not be summary data calculated for these parameters. These numbers of measurements are required to ensure that the results are indicative of overall summer conditions.

If you enrolled in Dissolved Oxygen/Temperature, the summary page will have a graph of one of the profiles taken during the late summer (typically August or September). If your lake stratifies, we will use a graph showing the earliest time of stratification, because identifying the timing of this condition and the depth at which it occurs is typically the most important use of dissolved oxygen measurements.

The back of the summary page will be an explanation of the Trophic Status Index and where your lake fits on that scale.

The rest of the report will be aquatic plant summaries, Score the Shore results, and larger graphs, including all Dissolved Oxygen/Temperature Profiles that you recorded. For Secchi Disk, Chlorophyll, and Phosphorus parameters, you need to have two years of data for a graph to make logical sense. Therefore if this is the first year you have enrolled in the CLMP, you will not receive a graph for these parameters.

Remember that some lakes see a lot of fluctuation in these parameters from year to year. Until you have eight years worth of data, consider all trends to be preliminary.

To learn more about the CLMP monitoring parameters or get definitions to unknown terms, check out the CLMP Manual, found at: https://micorps.net/wp-content/uploads/2021/03/CLMP-Manual-2019update2\_2021.pdf

## Thank you!

The CLMP leadership team would like to thank you for all of your efforts over the past year. The CLMP would not exist without dedicated and hardworking volunteers!

The CLMP Leadership Team is made of: Jo Latimore, Erick Elgin, Jean Roth, Tamara Lipsey, Mike Gallagher, Melissa DeSimone, and Paul Steen

## **Questions?**


If you have questions on this report or believe that the tabulated data for your lake in this report are in error please contact:

Paul Steen (psteen@hrwc.org), CLMP Data Analyst

# Second Fortune Lake, Iron County 2023 CLMP Results



## Secchi Disk Transparency (feet)



## Spring Phosphorus (parts per billion)

Year	# Samples	Min	Max	Average	Std. Dev	
2023	1	6.0	6.0	6.0	NA	
2018-2022	3	<=3 W	9.0	6.3	3.1	
2017	1	<5 T	<5 T	<5 T	NA	
2023 All CLMP Lakes	220	<5	220.0	20.7	21.3	
Spring Total Phosphorus (ppb) D p c 21			•		<b>.</b>	₽ =

## **Dissolved Oxygen and Temperature Profile**

2018

2016

This lake does not have recent (within 5 years) dissolved oxygen/water temperature data available. Consider enrolling in this parameter next year. Fish, insects, mollusks, and crustaceans need dissolved oxygen to live in water. By late summer, many lakes stratify, with cold anoxic water on the bottom and warm, oxygen rich water on the surface. Anoxic (oxygen-depleted) water occurring too close to the surface is a sign of nutrient enrichment. Understanding the pattern of dissolved oxygen and water temperature in a lake is important for assessing nutrient problems as well as the health of the biological community.

2020


2022

## Chlorophyll-a (parts per billion)

Second Fortune Lake does not have Chlorophyll-a data available. Consider enrolling in this parameter next year. Chlorophyll-a is the green photosynthetic pigment in the cells of plants. The amount of algae in a lake can be estimated by measuring the chlorophyll-a concentration in the water. As an algal productivity indicator, chlorophyll-a is used to determine the trophic status of a lake.

## Summer Phosphorus (parts per billion)

Year	# Samples	Min	Max	Average	Std. Dev	Carlson TSI	
2023	1	9.0	9.0	9.0	NA	36	
2018-2022	3	<5 T	7.0	5.3	1.5	29	
2017	1	5.0	5.0	5.0	NA	27	
2023 All CLMP Lakes	234	<= 3	150.0	17 4	15.3	45	



#### Summary

Average TSI	2023	2018-2022	2017
Second Fortune Lake All CLMP	36	31	29
Lakes	44	41	38

With a TSI score of 36 based on 2023 summer total phosphorus, this lake is rated between the oligotrophic and mesotrophic classification. The lake leans slightly more oligotrophic than mesotrophic.

There is too little data to assess long term trends. CLMP recommends eight years of consistent monitoring in order to develop a strong data baseline.

Reminder: 8 Secchi measurements are required in order to use the data in graphs and trends.

<sup>\* =</sup> Minimum # samples not met for average/median/TSI value

<sup>&</sup>lt;1.0 = Chlorophyll-a: Sample value is less than limit of quantification (<1 ppb).

W= Value is less than the detection limit (<3 ppb) T = Value reported is less than the reporting limit (5 ppb)

## **Trophic Status Index Explained**

In 1977, limnologist Dr. Robert Carlson developed a numerical scale (0-100) where the numbers indicate the level of nutrient enrichment. Using the proper equations, we can convert results from Summer Total Phosphorus, Secchi Depth, and Chlorophyll-a to this Trophic Status Index (TSI). The TSI numbers are furthermore grouped into general categories (oligotrophic, mesotrophic, eutrophic, and hypereutrophic), to quickly give us a way to understand the general nutrient level of any lake.

The tables below give the results-to-TSI conversions for the water quality data ranges normally seen in the CLMP. The formulas for this conversion can be found in the CLMP manual (link is on page 2 of this report).

Phosphorus		Secc	hi Depth		Chl	orophyll-a	
	<mark>Value</mark>		(ft)	TSI Value		(ppb)	TSI Value
<5	<27		>30	<28		<1	<31
6	30		25	31		2	37
8	34		20	34		3	41
10	37		15	38		4	44
12	40		12	42		6	48
15	43		10	44		8	51
18	46		7.5	48		12	55
21	48		6	52		16	58
24	50		4	57		22	61
32	54		<3	>61		>22	>61
36	56						
42	58						
48	60	TSI for	Second Fortu	ine Lake in i	<mark>2023</mark>		
>50	>61	Averag					
		Secchi	Disk				
		Summe					
		Chlorop	ohyll-a				
Oligotrophic	Oligo/Meso	Mesotrophic	Meso/Eutro	Eutrophi	с	Hypereutro	ohic
<36	36-40	41-45	46-50	51-61		>61	
0 5		9	5	0	55	0	5
<b>~</b> ~	,	4	4		- <b>P</b>	Ψ	4
C. P. T. T. T.	1111	L L L	1 1 1 1	6 6 6	11	1 9 9 10 1	TTE
	^ Average						

^ Total Phosphorus

**Oligotrophic:** Generally deep and clear lakes with little aquatic plant or algae growth. These lakes maintain sufficient dissolved oxygen in the cool, deep-bottom waters during late summer to support cold water fish, such as trout and whitefish.

Mesotrophic: Lakes that fall between oligotrophic and eutrophic. Mid-ranged amounts of nutrients.

**Eutrophic:** Highly productive eutrophic lakes are generally shallow, turbid, and support abundant aquatic plant growth. In deep eutrophic lakes, the cool bottom waters usually contain little or no dissolved oxygen. Therefore, these lakes can only support warm water fish, such as bass and pike.

**Hypereutrophic:** A specialized category of euthrophic lakes. These lakes exhibit extremely high productivity, such as nuisance algae and weed growth.

Site ID: 360177

## **Second Fortune, Iron County 2021 Exotic Aquatic Plant Watch Results**



The Exotic Aquatic Plant Watch was conducted on Second Fortune Lake in 2021.

This survey involves sampling at multiple locations around the lake to detect new invaders, and document the extent of known invaders. While notes on other plant species may be recorded during the survey, the effort focuses on five highly invasive species: Eurasian watermilfoil (Myriophyllum spicatum), starry stonewort (Nitellopsis obtusa), curly-leaf pondweed (Potamogeton crispus), European Frogbit (Hydrocharis morsusranae), and Hydrilla (Hydrilla verticillata).

The table below summarizes the results of the 2021 Exotic Aquatic Plant Watch.

## Second Fortune Lake, Iron County

#### **2021 Exotic Aquatic Plant Watch Results**

<u>Species</u>	<u>Status</u>	<u>Comments</u>	
Eurasian watermilfoil	not found		
Starry stonewort	not found		
Curly-leaf pondweed	not found		
European Frogbit	not found		
Hydrilla	not found		

#### . . . . . . .


Visit the MiCorps Data Exchange (https://micorps.net) or contact the lead volunteer on your lake for more details on the survey, including sampling locations, maps, and abundance information, and for information on past surveys.

## Second Fortune Lake, Iron County 2017 Score the Shore Results



The Score the Shore Habitat Assessment was conducted on Second Fortune Lake in 2017.

This assessment involves rating 1000 foot sections of shoreline for aquatic vegetation, shoreline vegetation, erosion, and erosion control practices (like sea walls). Each shoreline section is given three scores ranging from 0-100 for the categories of Littoral, Riparian, and Erosion Management. The three scores are averaged to produce a average section score. Then a total score is given to the entire lake by averaging all of the average section scores. A score of 0 indicates a shoreline that has been extremely disturbed by human impacts and no natural shoreline remains. A score of 100 indicates a shoreline that is nearly pristine.



#### How does your lake compare to others in the program?


Overall, the lakeshore habitat of Second Fortune Lake is doing well and scored higher than average when compared to other lakes in the program. All of the 1000 foot sections scored either Fair or Good: 1 fair, and 9 good.

The lake sections scored high for erosion control meaning that there are a low amount of sea walls, rock rip-rap, and other shoreline erosion structures, and scored high in the riparian score, meaning that (in general) there were plentiful non-mowed areas providing a buffer between houses and roads and the lake.


The littoral zone was the weak point in Second Fortune Lake's habitat (scoring an average of 77). A score of 77 is good, but if residents of want to improve the overall shoreline quality, this is the component to concentrate on. To improve the littoral zone score, leave woody debris in place and allow native aquatic vegetation to grow in the shallow waters.

Second Fortune Lake:				
Number of Sections:	10			
Number of Structures:	53			
Structure Density:	5.3			
Final Score:	89			


All 31 Participating Lakes from 2015-2017:				
Avg. Number of Sections:	16			
Avg. Number of Structures:	233			
Avg. Structure Density:	14.6			
Avg. Final Score:	66			



#### COOPERATIVE LAKES MONITORING PROGRAM SUMMER MEAN TRANSPARENCY



#### COOPERATIVE LAKES MONITORING PROGRAM SPRING TOTAL PHOSPHORUS



#### COOPERATIVE LAKES MONITORING PROGRAM SUMMER TOTAL PHOSPHORUS

